


# **DETERMINATION OF MYCOTOXINES IN MEAT**

#### **PROTOCOL OF PURIFICATION**

#### Sample preparation

25g of meat are mixed during 2 minutes with 100mL of a solution of deionized water/ acetonitrile (50/50). Then the mixture is filtered through filter paper (4-7 $\mu$ m). This solution was then diluted 10 times with deionized water to obtain the loading solution used for the clean-up protocol

Loading



# AFFINIMIP<sup>®</sup> SPE DEOXYNIVALENOL

Purification with a 6mL/100mg AFFINIMIP<sup>®</sup> SPE DEOXYNIVALENOL cartridge

#### Equilibration

- •2mL Acetonitrile
- 2mL Water

#### Loading solution

Up to 6mL of loading solution

#### Washing of interferences

•3mL NaHCO<sub>3</sub> 1% solution

## Drying 30s

Washing of interferences

1mL Diethyl Ether


#### Elution (E)

4mL Ethyl Acetate

The elution fraction was then concentrated and diluted to mobile phase before HPLC analysis.

# **Recovery for DON > 90%\***

\*Tested at 8000µg/kg



UV Chromatograms (220nm) obtained after clean-up of MEAT with AFFINIMIP® SPE DEOXYNIVALENOL.

- Blue trace for spiked with 8000µg/kg of DEOXYNIVALENOL
- Red trace for the blank sample

#### HPLC Method with LC-MS

HPLC Column: Hypersil gold column (50mm x 2.1mm) Mobile phase: 95/5 0,1 Formic acid in water / Acetonitrile Flow rate: 0.2mL/min Injection volume: 20μL.

**Ordering information - Catalog number:** FS117-02B for 25 cartridges, 6mL FS117-03B for 50 cartridges, 6mL

## **AFFINIMIP<sup>®</sup> SPE ZEARALENONE**

# Purification with a 3mL/100mg AFFINIMIP® SPE ZEARALENONE cartridge

#### Equilibration

•2mL Acetonitrile

Elution

•2mL Water

#### Loading solution

Up to 6mL of loading solution

#### Washing of interferences

•6mL 40/60 Acetonitrile/deionized Water Drying 3-5min

#### **Elution (E)**

2mL 2/98 Acetic Acid/Methanol The elution fraction was then concentrated and diluted to mobile phase before HPLC analysis.

## Recovery for ZON > 80%\* \* Tested at 960µg/kg

# Zearalenone Zearalenone

SIM Chromatograms obtained after clean-up of MEAT with AFFINIMIP<sup>®</sup> SPE ZEARALENONE.

- Blue trace for spiked with 960µg/kg of ZEARALENONE
- Red trace for the blank sample

#### HPLC Method with LC-MS

HPLC Column: Hypersil gold column (50mm x 2.1mm) Mobile phase: 73/27 0,1 Formic acid in water / Acetonitrile Flow rate: 0.2mL/min Injection volume: 20µL.

Ordering information - Catalog number: FS100-02 for 25 cartridges, 3mL FS100-03 for 50 cartridges, 3mL